REPUBLIC OF LEBANON MINISTRY OF PUBLIC HEALTH Lebanon Clinical Trials Registry

Safety Study of Crushed Deferasirox Film Coated Tablets in Pediatric Patients With Transfusional Hemosiderosis (MIMAS)

24/08/2025 04:42:26

Primary registry identifying number	Protocol number
LBCTR2019030206	CICL670F2429
MOH registration number	
32772/2018	
Study registered at the country of origin	Study registered at the country of origin: Specify
Yes	
Type of registration	Type of registration: Justify
Retrospective	LCTR was recently initiated, original file was previously submitted by Paper
Date of registration in national regulatory agency 01/08/2018	
Primary sponsor	Primary sponsor: Country of origin
Novartis Pharma Services Inc.	Novartis Pharmaceuticals
Date of registration in primary registry	Date of registration in national regulatory agency
12/03/2019	01/08/2018
Public title	Acronym
Safety Study of Crushed Deferasirox Film Coated Tablets in Pediatric Patients With Transfusional Hemosiderosis (MIMAS)	
Scientific title	Acronym
A Single-arm Interventional Phase IV, Post-authorisation Study Evaluating the Safety of Pediatric Patients With Transfusional Hemosiderosis Treated With Deferasirox Crushed Film Coated Tablets	
Brief summary of the study: English	
The study employs an interventional, prospective, single arm, open label, global, multicenter,non-randomized trial design to monitor and assess the safety profile of the crushed deferasirox FCT in pediatric patients between age ≥2 to <6 with transfusional hemosiderosis over 24 weeks. This study will aim to enroll at least 40 patients.	
Brief summary of the study: Arabic	
الرابعة بعد الترخيص لتقييم سلامة المرضى الاطفال المصابين بالحدد الدموي (الهيموسيدروسز) ذي الصلة بنقل الدم والمعالجين بأقراص ديفيرازيروكس المسحوقة المغلفة بطبقة رقيقة	دراسة تدخلية وحيدة المجموعة في المرحلة
Health conditions/problem studied: Specify	
Transfusional Hemosiderosis	
Interventions: Specify	
Drug: Deferasirox Deferosirox is provided in tablet forms of 90, 180 and 360mg. Tablets rr	nust be crushed.
Other Name: ICL670	

1.Patients ≥2 to <6 years old diagnosed with transfusional hemosiderosis

 \sim

REPUBLIC OF LEBANON Lebanon Clinical Trials Registry

2.Documented history of red blood cell transfusions

3.Written informed consent/assent before any study-specific procedures. The consent will be obtained from caregiver(s) or patient's legal representative. Investigators will also obtain assent of patients according to local, regional, or national regulations. 4.For patients on prior DFX: Serum ferritin (SF) >500 ng/mL, measured at screening visit 1 and requiring a DFX daily dose equivalent to FCT ≥ 7mg/kg/day.

5.For patients on a prior chelator other than DFX (e.g. deferiprone or deferoxamine) or chelation naive: Serum ferritin (SF) >1000 ng/mL measured at screening visits 1 and 2.

Key inclusion and exclusion criteria: Gender	Key inclusion and exclusion criteria: Specify gender
Both	
Key inclusion and exclusion criteria: Age minimum	Key inclusion and exclusion criteria: Age maximum
2	6

Key inclusion and exclusion criteria: Exclusion criteria

1.Patients that receive more than one iron chelator at the same time as current iron chelation treatment. (Patients who have received combination therapy in their medical history but are currently being treated with a single ICT agent are eligible.)

2.Patients continuing on deferoxamine or deferiprone in addition to study treatment.

(Patients switching to or continuing on deferasirox are eligible).

3. Unresolved adverse events if the patient was previously treated with deferiprone or deferoxamine or deferasirox.

4. Significant proteinuria as indicated by a urinary protein/creatinine ratio > 0.5 mg/mg in a non-first void sample urine measured at screening visit 1.

5.Serum creatinine > age adjusted ULN measured at any screening visit

6.Creatinine clearance below 90 mL/minute measured at any screening visit. Creatinine clearance using the Schwartz formula will be estimated from serum creatinine measured at each respective visit.

7.ALT and/or AST > 2.5 x ULN measured at screening visit 1.

8.Total bilirubin (TBIL) >1.5 x ULN measured at screening visit 1.

9.Patients with significant impaired GI function or GI disease that may significantly alter the absorption of oral deferasirox FCT (e.g. ulcerative diseases, uncontrolled nausea, vomiting, diarrhea, malabsorption syndrome, or small bowel resection).

10. History of and/or laboratory evidence of active Hepatitis B or Hepatitis C (HBsAg in the absence of HBsAb OR HCV Ab positive with HCV RNA positive.

11.Liver disease with severity of Child-Pugh Class B or C.

12. History of hypersensitivity to any of the study drug or excipients.

13.Patients participating in another clinical trial or receiving an investigational drug.

14.Patients with a known history of HIV seropositivity.

15.Patients unwilling or unable to comply with the protocol.

16. History of malignancy of any organ system, treated or untreated, within the past 5 years whether or not there is evidence of local recurrence or metastases, with the exception of localized basal cell carcinoma of the skin.

17. Significant medical condition interfering with the ability to partake in this study (e.g.

uncontrolled hypertension, unstable cardiac disease not controlled by standard medical therapy, systemic disease: cardiovascular, renal, hepatic, etc.).

18.Female patients who reach menarche and they or their caregivers refuse pregnancy testing and/or if there is a positive pregnancy test result.

Type of study

Interventional

Type of intervention	Type of intervention: Specify type
Pharmaceutical	N/A
Trial scope	Trial scope: Specify scope
Therapy	N/A
Study design: Allocation	Study design: Masking
N/A: Single arm study	Open (masking not used)
Study design: Control	Study phase
N/A	4
Study design: Purpose	Study design: Specify purpose

REPUBLIC OF LEBANON Lebanon Clinical Trials Registry

Treatment	N/A	
Study design: Assignment Single	Study design: Specify assignment N/A	
IMP has market authorization	IMP has market authorization:	Specify
Yes, Lebanon and Worldwide	USA, UK, France, Germany, Net Italy	
Name of IMP	Year of authorization	Month of authorization
Deferasirox Film Coated Tablets "Jadenu"	2017	10
Type of IMP Others		
Pharmaceutical class Non-chiral, Tridentate ligand iron chelator		
Therapeutic indication Iron Over Load		
Therapeutic benefit Treatment of Iron Overload symptoms		
Study model N/A	Study model: Explain model N/A	
Study model: Specify model N/A		
Time perspective N/A	Time perspective: Explain time	e perspective
Time perspective: Specify perspective N/A		
Target follow-up duration	Target follow-up duration: Unit	t
Number of groups/cohorts		
Bissessimon retention	Biognosimon des suisties	
Biospecimen retention None retained	Biospecimen description Local Lab tests at site	
Target sample size 3	Actual enrollment target size	
Date of first enrollment: Type	Date of first enrollment: Date	

panon Clinical Trials Registry

Actual	10/10/2018
Date of study closure: Type	Date of study closure: Date
Actual	11/03/2020
Recruitment status	Recruitment status: Specify
Recruiting	
Date of completion	
22/04/2019	
IPD sharing statement plan	IPD sharing statement description
No	undecided
Additional data URL	
https://clinicaltrials.gov/ct2/show/record/NCT03372083?id=CIC	L670F2429&rank=1&view=record
Admin comments	
Trial status	

Approved

Secondary Identifying Numbers		
Full name of issuing authority	Secondary identifying number	
Clinical Trials. gov	NCT03372083	

Sources of Monetary or Material Support
Name
Novartis Pharma Services Inc.
Name

Secondary Sponsors	
Name	
NA	



Contact for Public/Scientific Queries						
Contact type	Contact full name	Address	Country	Telephone	Email	Affiliation
Public	Ali Taher	Beirut	Lebanon	009613755 669	ataher@aub.edu. lb	Chronic Care Center
Scientific	Hind Khairallah	Sin El Fil	Lebanon	+961 1 512002 Ext. 271	Hind.Khairallah@ fattal.com.lb	Khalil Fattal et Fils s.a.l.

Centers/Hospitals Involved in the Study				
Center/Hospital name Name of principles investigator		Principles investigator speciality	Ethical approval	
Chronic Care Center	Dr Ali Taher	Hematology	Approved	

Ethics Review					
Ethics approval obtained Approval date Contact name		Contact name	Contact email	Contact phone	
Chronic Care Center	26/07/2018	Michele Abi saad	cccmas@chroniccare.org.lb	+961 3 664 310	
American University of Beirut Medical Center	05/07/2018	Fuad Ziyadeh	fz05@aub.edu.lb	+961 (0) 1 350 000 ext:5445	

Countries of Recruitment
Name
Egypt
Lebanon
Italy
Oman
Russian Federation
Saudi Arabia
Thailand
United Arab Emirates



Health Conditions or Problems Studied				
Condition	Code	Keyword		
Transfusional Hemosiderosis	Thalassaemia, unspecified (D56.9)	Transfusional Hemosiderosis		

Interventions				
Intervention	Description	Keyword		
Physical examination, Vitals, Height, Weight, Hematology, Chemistry, urinalisis, ECG, Ocular assessment, Auditory assessment	Physical examination, Vitals, Height, Weight, Hematology, Chemistry, urinalisis, ECG, Ocular assessment, Auditory assessment	ICF, IMP, Lab tests , diary completion		

Primary Outcomes				
Name	Time Points	Measure		
Percentage of patients with selected gastrointestinal disorders	24 weeks	24 wks		
To assess the safety of crushed deferasirox FCT with respect to selected gastrointestinal (GI) disorders	through out the study	through out the study		

Key Secondary Outcomes			
Name	Time Points	Measure	
 Percentage of patients who experienced AEs suspected to be related to study drug 	24 weeks	24 wks	
•Change from baseline ECGs up	24 weeks	24 weeks	
•Change from baseline serum ferritin (SF)	24 weeks	24 weeks	
•Absolute change for serum creatinine	24 weeks	24 weeks	
•Absolute change for creatinine clearance UPCR	24 weeks	24 weeks	
Palatability Questionnaire Score	24 weeks	24 weeks	



Trial Results Summary results Study results globally Date of posting of results summaries Date of posting of results summaries Date of first journal publication of results Results URL link Baseline characteristics Participant flow Adverse events Outcome measures URL to protocol files