

### A NON-INTERVENTIONAL, MULTICENTER, MULTIPLE COHORT STUDY INVESTIGATING THE OUTCOMES AND SAFETY OF ATEZOLIZUMAB UNDER REAL-WORLD CONDITIONS IN PATIENTS TREATED IN ROUTINE CLINICAL **PRACTICE**

20/08/2025 08:55:44

-					4 8	
N	ЛЭІ	$\mathbf{n}$		ma	177	חר
IIV.	11-11		 			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

Primary registry identifying number

LBCTR2020033423

MOH registration number

Study registered at the country of origin

Type of registration

Prospective

Date of registration in national regulatory agency

07/01/2020

**Primary sponsor** 

F. HOFFMANN-LA ROCHE LTD

Date of registration in primary registry

20/03/2020

**Public title** 

A NON-INTERVENTIONAL, MULTICENTER, MULTIPLE COHORT STUDY INVESTIGATING THE OUTCOMES AND SAFETY OF ATEZOLIZUMAB UNDER REAL-WORLD CONDITIONS IN PATIENTS TREATED IN ROUTINE CLINICAL PRACTICE

Scientific title

A NON-INTERVENTIONAL, MULTICENTER, MULTIPLE COHORT STUDY INVESTIGATING THE OUTCOMES AND SAFETY OF ATEZOLIZUMAB UNDER REAL-WORLD CONDITIONS IN PATIENTS TREATED IN ROUTINE CLINICAL PRACTICE

Brief summary of the study: English

Protocol number

MO40653

Study registered at the country of origin: Specify

Type of registration: Justify

N/A

Primary sponsor: Country of origin

Germany

Date of registration in national regulatory agency

07/01/2020

Acronym

**IMReal** 

Acronym

**IMReal** 



In the real world, the clinical benefit, and safety of atezolizumab have not yet been established. This study will contribute to the understanding of the short and long-term outcomes and safety of atezolizumab in patients with locally advanced/metastatic UC, locally advanced/metastatic NSCLC and ES-SCLC in the real-world setting.

The study aims to address the following research questions in patients treated with atezolizumab for the first time for the approved indications, in the real-world setting of routine clinical practice:

- •What are the clinical outcomes of atezolizumab?
- •What is the quality of life of patients treated with atezolizumab?
- •What is the safety profile of atezolizumab?
- •What is the sociodemographic and clinical profile of patients treated with atezolizumab?
- •What are the characteristics of atezolizumab treatment (e.g. treatment duration,
- •discontinuation, interruption, number of cycles)?
- •What are the characteristics of prior and subsequent cancerrelated therapies?
- •What is the healthcare resource utilisation and associated cost of atezolizumab treatment?
- •What are non-healthcare consequences and associated costs of atezolizumab?

#### Brief summary of the study: Arabic

ان الهدف من هذه الدراسة الرصدية معرفة كيف يعمل العلاج باتيزوليزوماب في "الحياة الحقيقية"، في سياق الممارسة الطبية الروتينية. كذلك، بواسطة هذه الدراسة، يمكن الاستحصال على معلومات اضافية تتعلق بالمأمونية، اي معلومات حول مدى تحمّل المريض للدواء عندما . يوصف له في اطار الممارسة الطبية اليومية . . يوصف له في اطار الممارسة الطبية اليومية . هناك العديد من العوامل التي لا يمكن دراستها الا في الممارسة اليومية، على وجه الخصوص اذا كان عدد المرضى المشاركين كبيراً على سبيل المثال، يمكن اكتشاف أثار جانبية غير معروفة كثيراً للدواء، كما يمكن ملاحظة تأثير تغيير الجرعات او الادوية المصاحبة، الخ. من شأن هذه المعلومات ان تساعد الاطباء ليعرفوا بصورة افضل كيفية استخدام اتيزوليزوماب، كما من شأنها ان تحسّن الى الحد الاقصى علاج المرضى

#### Health conditions/problem studied: Specify

Patients diagnosed with locally advanced/metastatic non-small cell lung cancer (NSCLC) after prior chemotherapy. Patients with epidermal growth factor receptor (EGFR) activating mutations or anaplastic lymphoma kinase (ALK)- positive tumor mutations should also have received targeted therapy

#### Interventions: Specify

.المشاركين

Non-interventional study.

#### Key inclusion and exclusion criteria: Inclusion criteria

Patients must meet the following criteria for study entry:

- 1.Patient must have one of the following confirmed diagnoses for which atezolizumab is locally approved in the SmPC:
- oAs monotherapy for the treatment of adult patients with locally advanced/metastatic UC after prior platinum-containing chemotherapy (Cohort 1 LOT2+ mUC).

oAs monotherapy for the treatment of adult patients with locally advanced/metastatic NSCLC after prior chemotherapy. Patients with EGFR activating mutations or ALK- positive tumour mutations should also have received targeted therapy (i.e. ALK /EGFR-TKIs) before receiving atezolizumab (Cohort 2 LOT2+ NSCLC).

oln combination with bevacizumab, paclitaxel and carboplatin for the first line treatment of adult patients with metastatic non-squamous NSCLC. Patients with EGFR activating mutations or ALK- positive tumour mutations should also have received targeted therapy (i.e. ALK /EGFR-TKIs). (Cohort 3 LOT1 NSCLC).

oln combination with carboplatin and etoposide, is indicated for the first-line treatment of adult patients with extensive-stage small cell lung cancer (ES-SCLC) (Cohort 4 LOT1 ES-SCLC)

- 2.Patient is prescribed atezolizumab therapy for the first time
- 3.Decision to prescribe atezolizumab must be made and documented prior to inclusion into the study and must follow local clinical practice.
- 4.Patient is aged ≥ 18 years or older at the index date.
- 5. Patient has signed an informed consent form according to local regulations.
- 6.Data collection can only start after the signing of inform consent and not more than 28 days after initiation of atezolizumab. In countries where enrollment is only allowed after the treatment start, enrollment must be preceded by the administration of the first cycle of atezolizumab.

100

Key inclusion and exclusion criteria: Gender

Key inclusion and exclusion criteria: Specify gender

Both

Key inclusion and exclusion criteria: Age minimum

Key inclusion and exclusion criteria: Age maximum

8

#### Key inclusion and exclusion criteria: Exclusion criteria

Patients who meet any of the following criteria will be excluded from study entry:

1. Patients not receiving treatment for a disease with atezolizumab according to standard of care and in line with the current summary of





product characteristics (SPC) or local labelling \*

- 2.Concomitant anti-cancer therapy at the time of starting atezolizumab on the index date, not part of locally approved combination therapy with atezolizumab.
- 3. Treatment with atezolizumab as part of a clinical trial or for compassionate use as part of an access or compassionate use program.
- 4. Patients not receiving atezolizumab, but a biosimilar or non-original biologic.

Type of study Observational

Type of intervention

N/A

Trial scope

N/A

Study design: Allocation

N/A

Study design: Control

N/A

Study design: Purpose

N/A

Study design: Assignment

N/A

IMP has market authorization

Name of IMP

Type of IMP

Pharmaceutical class

Therapeutic indication

Therapeutic benefit

Study model

Cohort

Study model: Specify model

N/A

Type of intervention: Specify type

N/A

Trial scope: Specify scope

N/A

Study design: Masking

N/A

Study phase

N/A

Study design: Specify purpose

N/A

Study design: Specify assignment

N/A

IMP has market authorization: Specify

Year of authorization Month of authorization

Study model: Explain model

The study will be split into separate cohorts based on the approved indications for atezolizumab treatment, excluding cisplatin ineligible patients receiving atezolizumab as first line of therapy (LOT1) for locally advanced/metastatic urothelial cancer (locally advanced/metastatic UC). The study may be amended for inclusion of new cohorts as these are approved in the participating countries. Patients will be included into each cohort based on their indication for receiving atezolizumab.

Lebanon is participating in Cohort 2 (NSCLC LOT2 plus later lines [LOT2+]): Patients diagnosed with locally advanced/metastatic non-small cell lung cancer (NSCLC) after prior chemotherapy. Patients with epidermal growth factor receptor (EGFR) activating mutations or anaplastic lymphoma kinase (ALK)- positive tumor mutations should also have received targeted therapy.



Time perspective

Prospective

Time perspective: Specify perspective

N/A

Target follow-up duration

5

Number of groups/cohorts

4

Biospecimen retention

None retained

Target sample size

10

Date of first enrollment: Type

Anticipated

Date of study closure: Type

Anticipated

Recruitment status

Pending

Date of completion

01/06/2025

IPD sharing statement plan

Yes

Time perspective: Explain time perspective

This is a non-interventional, multi-country, multi-centre, multiple cohort prospective study, with retrospective collection of prior medical/treatment history data from medical records, designed to assess the real-world outcomes and safety of atezolizumab for indications in the existing label in the real-world setting of routine clinical practice. The study will collect data before, during and after treatment with atezolizumab.

Target follow-up duration: Unit

years

Biospecimen description

Non retained

Actual enrollment target size

Date of first enrollment: Date

01/04/2020

Date of study closure: Date

01/06/2025

**Recruitment status: Specify** 

IPD sharing statement description



During this study, health and personal information about subjects will be collected. This section describes the protection, use, and sharing of information, which consists of the following:

- · Information in the medical record, which is held by Sites.
- Information that is collected or produced during this study ("study data"), which is held by sites, Roche, Roche affiliates, and Roche's representatives.

Subject privacy is very important, and Roche uses many safeguards to protect privacy, in accordance with applicable data privacy laws and laws related to the conduct of clinical trials. Subject study data will be labelled with a patient identification (ID) number that is unique and not related to or derived from information that identifies subject (such as name, picture, or any other personally identifying information). Roche, Roche affiliates, and Roche's representatives will only have access to study data and samples labelled with a patient ID number, except as described below. Subjects medical record, which includes personal information that can identify subjects, will not be accessed for the purposes of this study, except as described below:

Information (which includes information in medical record that can identify subjects) may need to be reviewed to make sure the study is being done properly or to check the quality of the information. This information will be kept private. The following people and groups of people may and/or copy this information:

• Study monitors of Roche and/or CRO, a company hired by Roche to perform certain study activities

- The Institutional Review Board or Ethics Committee
- · Regulatory authorities

Roche, Roche affiliates, and Roche's collaborators and licensees (people and companies who partner with Roche) may use study data labelled with patient ID number for research purposes or to advance science and public health.

Study data may be submitted to government or other health research databases or shared with researchers, government agencies, companies, or other groups that are not participating in this study. These data may be combined with or linked to other data and used for research purposes, to advance science and public health, or for analysis, development, and commercialization of products to treat and diagnose disease. These data will not include information that identifies subjects, and extra steps will be taken to safeguard privacy.

Subject information will not be given to insurance company or employer, unless required by law. If the results from this study are published in a medical journal or presented at a scientific meeting, subjects will not be identified.

Information from this study will be retained by Sites for 15 years after the end of the study. In addition, Roche will retain the study data for up to 25 years after the end of the study.

Additional data URL

Admin comments

Trial status

Approved

Secondary Identifying Numbers		
Full name of issuing authority	Secondary identifying number	
NA	NA	



#### **Sources of Monetary or Material Support**

Name

F. HOFFMANN-LA ROCHE LTD

### **Secondary Sponsors**

Name

NA

Contac	Contact for Public/Scientific Queries						
Contact type	Contact full name	Address	Country	Telephone	Email	Affiliation	
Public	Dani Abi Gerges	Middle East Institute of Health - University Hospita, Bsalim, Metn P.O. Box 60-387 Metn, Lebanon	Lebanon	009614712 111	abgerges@idm.n et.lb	Middle East Institute of Health - University Hospital	
Scientific	Dani Abi Gerges	Middle East Institute of Health - University Hospita, Bsalim, Metn P.O. Box 60-387 Metn, Lebanon	Lebanon	009614712 111	abgerges@idm.n et.lb	Middle East Institute of Health - University Hospital	

Centers/Hospitals Involved in the Study				
Center/Hospital name	Name of principles investigator	Principles investigator speciality	Ethical approval	
Middle East Institute of Health Hospital	Dr. Dani Abigerges	Hemato-Oncology	Approved	

Ethics Review				
Ethics approval obtained	Approval date	Contact name	Contact email	Contact phone
Middle East Institute of Health	07/01/2020	Dr. Rania Haddad	r_haddad@meih.org	009614712111 ext. 6063



Countries of Recruitment
Name
Hungary
Argentina
France
Spain
Bulgaria
Austria
Belgium
United Kingdom
Netherlands
Lithuania
Italy
Croatia
Portugal
Poland
Slovenia
Lebanon

Health Conditions or Problems Studied			
Condition	Code	Keyword	
Non Small Cell Lung Cancer	Other disorders of lung (J98.4)	NSCLC	

Interventions			
Intervention	Description	Keyword	
Non-interventional study	NA	NA	



Primary Outcomes				
Name	Time Points	Measure		
•To estimate overall survival (OS) at two years and at the end of the study.	Overall Survival - OS	Time from index date until date of death from any cause		
•To estimate overall survival (OS) at two years and at the end of the study.	OS at 2 years	Proportion of patients alive 2 years after the index date		

Key Secondary Outcomes				
Name	Time Points	Measure		
To evaluate the clinical benefit of atezolizumab	Time to loss of clinical benefit	Time from the index date to loss of clinical benefit as assessed by the treating physician (single or multiple reasons possible: e.g. disease progression, deterioration in ECOG performance status, patient preference, death, etc.). Clinical benefit is defined as:  • Absence of symptoms and signs indicating unequivocal progression of disease • No decline in ECOG performance status • Absence of tumour progression at critical anatomical sites that cannot be readily managed and stabilized by protocol-allowed medical interventions prior to repeat dosing • Evidence of clinical benefit as assessed by the physician according to local clinical standard of care		
To estimate PFS	Progression-free survival - PFS	Time from index date to death or disease progression (DP; physician assessed according to local clinical standard of care or as per RECIST if available)		
To estimate the ORR, DCR and DoR	Objective response rate - ORR	ORR is the proportion of patients who have a BOR of CR or PR. Best overall response (BOR) for each patient is the best response (physician assessed according to local clinical standard of care or as per RECIST if available) achieved after the index date prior to initiation of any subsequent treatment.		
To estimate the ORR, DCR and DoR	Time to response	Time from index date to first objective tumour response, CR or PR (physician assessed according to local clinical standard of care or as per RECIST if available)		
To estimate the ORR, DCR and DoR	Duration of response - DoR	Time from first documentation of CR or PR (whichever occurs first) after index until death or PD (physician assessed according to local clinical standard of care or as per RECIST if available)		
To estimate the ORR, DCR and DoR	Disease control rate - DCR	Percentage of patients who have achieved CR, PR and stable disease (physician assessed according to local clinical standard of care or as per RECIST if available) at least 12 weeks after the index date		
To estimate the ORR, DCR and DoR	Duration of DCR	Time from first documentation of CR, PR or stable disease (whichever occurs first) after index until death or PD.		
To evaluate HRQoL	The EQ-5D-5L during and after atezolizumab treatment will be used to assess HRQoL	The EQ-5D-5L is a standardized instrument for measuring generic health status and has been used in the atezolizumab clinical trials. It includes two components, health state description and evaluation.		
To describe baseline patient characteristics relevant to treatment and/or prognosis	Refer to protocol page 44 for table of patient characteristics	To describe baseline patient characteristics relevant to treatment and/or prognosis		
To describe the characteristics of atezolizumab treatment, prior and subsequent cancer related therapies and estimate time to initiation of subsequent LOTs	Refer to protocol page 45,46,47 for Characteristics of atezolizumab treatment, and prior/subsequent cancer-related therapies	The time from diagnosis to the index date and time from last LOT to the index date will also be calculated.		



Trial Results	
Summary results	
Study results globally	
Date of posting of results summaries	Date of first journal publication of results
Results URL link	
Baseline characteristics	
Participant flow	
Adverse events	
Outcome measures	
URL to protocol files	