REPUBLIC OF LEBANON Lebanon Clinical Trials Registry

A Randomized, Double-blind, Placebo-controlled, Multicenter Study to Assess the Safety and Efficacy of Inclacumab in Participants with Sickle Cell Disease Experiencing Vasoocclusive Crises

20/08/2025 12:44:03 Main Information Primary registry identifying number Protocol number LBCTR2021064791 GBT2104-131 MOH registration number Study registered at the country of origin Study registered at the country of origin: Specify No Study is currently under registration in the US Type of registration Type of registration: Justify Prospective N/A Date of registration in national regulatory agency 18/06/2021 **Primary sponsor** Primary sponsor: Country of origin Global Blood Therapeutics, Inc. USA Date of registration in primary registry Date of registration in national regulatory agency 13/07/2021 18/06/2021 Public title Acronym A Randomized, Double-blind, Placebo-controlled, Multicenter Study to Assess the Safety and Efficacy of Inclacumab in Participants with Sickle Cell Disease Experiencing Vaso-occlusive Crises Scientific title Acronym A Randomized, Double-blind, Placebo-controlled, Multicenter Study to Assess the Safety and Efficacy of Inclacumab in Participants with Sickle Cell Disease Experiencing Vaso-occlusive Crises Brief summary of the study: English This is a Phase 3, randomized, double-blind, placebo-controlled, 2arm, multi-center, parallel-group study. The primary objective of this study is to evaluate the safety and efficacy of treatment every 12 weeks with inclacumab, a P-selectin inhibitor, to reduce the incidence of vaso-occlusive crises (VOCs) in participants with sickle cell disease (SCD). Additional objectives of the study are to evaluate the pharmacokinetics (PK) and pharmacodynamics (PD) of inclacumab, the presence of anti-drug antibodies (ADAs), and changes in quality of life (QOL).

Brief summary of the study: Arabic

Bhei summary of the study. Arabi

REPUBLIC OF LEBANON MINISTRY OF PUBLIC HEALTH Lebanon Clinical Trials Registry

هذه هي دراسة في المرحلة الثالثة متعددة المراكز، عشوائية التوزيع، مزدوجة التعمية، مراقبة بدواء وهمي

أسبو تحا باستعمال عقار إنكالكوماب لتقليل حدوث نوبات انسداد اللوعية الدموية12الهدف الرئيس لهذه الدراسة هو تقييم أمان وفعالية العالج كل (PD) والديناميكا الدوائية (PK) لدى المشاركين المصابين بمرض فقر الدم المنجلي. تتمثل األهداف الإضافية للدراسة في تقييم الحرائك الدوائية . (QD) والديناميكا الدوائية (PK) لدى المشاركين المصابين بمرض فقر الدم المنجلي. تتمثل األهداف الإضافية للدراسة في تقييم الحرائك الدوائية

Health conditions/problem studied: Specify

Patients with a diagnosis of sickle cell disease who have experienced between 2 and 10 vaso-occlusive crises in the 12 months preceding enrollment in this study.

Interventions: Specify

This study will assess the safety and efficacy of inclacumab in reducing the frequency of VOCs in approximately 240 adult and adolescent participants (\geq 12 years of age) with SCD. Initial enrollment will include participants \geq 16 years of age until the independent Data Monitoring Committee (DMC) determines that adequate safety and PK data support the enrollment of participants 12 to 15 years of age. Eligible participants will be randomized with a 1:1 ratio into one of two treatment arms as follows:

- Inclacumab 30 mg/kg administered IV Q12W; or

- Placebo administered IV Q12W.

At the time of randomization, participants will be stratified by Baseline hydroxyurea (HU) use (yes; no), number of VOCs (2 to 4; 5 to 10) in the preceding 12 months, and geographic region (North America; sub-Saharan Africa; Europe/rest of world).

The total duration of treatment for each participant will be 48 weeks. Doses will be administered at Day 1 and Weeks 12, 24, and 36 with blood levels expected to be in the target range through at least Week 48.

Key inclusion and exclusion criteria: Inclusion criteria

Inclusion Criteria:

1. Participant has a confirmed diagnosis of SCD (HbSS, HbSC, HbSβ0

thalassemia, or HbSβ+ thalassemia genotype).

Documentation of SCD genotype is required and may be based on documented history of laboratory testing or confirmed by laboratory testing during Screening.

2. Participant is male or female, ≥ 12 years of age at the time of informed

consent.

NOTE: Initial study enrollment will include participants ≥ 16 years of age until the DMC determines that adequate safety and PK data support the

enrollment of participants 12 to 15 years of age. Sites will be informed by the Sponsor when participants 12 to 15 years of age may be enrolled. 3. Participant has experienced between 2 and 10 VOCs within the 12 months prior to the Screening Visit as determined by documented medical history.

A prior VOC is defined as an acute episode of pain which:

- Has no medically determined cause other than a vaso-occlusive event,

and

- Results in a visit to a medical facility (hospital, emergency department, urgent care center, outpatient clinic, or infusion center) or results in a remote contact with a healthcare provider; and

- Requires parenteral narcotic agents, parenteral nonsteroidal anti-inflammatory drugs (NSAIDs), or an increase in treatment with oral narcotics.

4. Participants receiving erythropoiesis-stimulating agents (ESA, erythropoietin [EPO]) must be on a stable dose for at least 90 days prior to the Screening Visit and expected to continue with the stabilized regimen throughout the course of the study.

5. Participants receiving HU, L-glutamine, or voxelotor (Oxbryta) must be on a stable dose for at least 30 days prior to the Screening Visit and

expected to

continue with the stabilized regimen throughout the course of the study.

6. Participant has adequate venous access, in the opinion of the Investigator, to comply with study procedures.

7. Participant understands the study procedures and agrees to participate in the study by giving written informed consent or parental permission/written assent.

8. Women of childbearing potential (WOCBP) are required to have a

negative serum pregnancy test at the Screening visit and negative urine

pregnancy test on all subsequent clinic visits and must agree to use a highly effective method of contraception throughout the study period and for at least 165 days after dosing.

Female participants will not be considered of childbearing potential if they are pre-menarchal, surgically sterile (hysterectomy, bilateral salpingectomy, tubal ligation, or bilateral oophorectomy) or postmenopausal (no menses for 12 months without an alternative medical

cause, confirmed by follicle-stimulating hormone test results).

Key inclusion and exclusion criteria: Gender	Key inclusion and exclusion criteria: Specify gender	
Both		
Key inclusion and exclusion criteria: Age minimum	Key inclusion and exclusion criteria: Age maximum	
12	90	
Key inclusion and exclusion criteria: Exclusion criteria		
Exclusion Criteria		

Candidates will be excluded from study entry if any of the following exclusion criteria exist at

REPUBLIC OF LEBANON

Lebanon Clinical Trials Registry

Screening or Baseline visits or at the timepoint specified in the individual criterion listed.

1. Participant is receiving regularly scheduled red blood cell (RBC) transfusion therapy (also termed chronic, prophylactic, or preventative transfusion).

2. Participant is taking or has received crizanlizumab (ADAKVEO) within 90 days prior to the Screening Visit.

3. Participant weighs > 133 kg (292 lbs.).

4. Participant has a significant active and poorly controlled (unstable) hepatic disorder clearly unrelated to SCD

5. Participant has any of the following laboratory values at screening:

a. Absolute neutrophil count (ANC) < 1.0 × 109/L

b. Platelet count < 80 × 109/L

c. Hemoglobin < 4.0 g/dL for adults and < 5.0 g/dL for participants ages 12 to < 18 years of age

d. Estimated glomerular filtration rate (eGFR) < 30 mL/min using Chronic Kidney Disease-Epidemiology Collaboration (CKD-EPI) formula in adults, and Schwartz formula in adolescents

6. Participant has known active (symptomatic) COVID infection or tests positive for COVID-19 during Screening.

7. Participant has a history of unstable or deteriorating cardiac or pulmonary disease within 6 months prior to consent including severe or

unstable pulmonary hypertension.

8. Participant has had treatment for a malignancy within the 12 months prior to the Screening Visit (except non-melanoma skin cancer and in situ cervical cancers).

9. Participant has had a stroke within the 2 years prior to the Screening Visit.

10. Participant has a positive test indicative of an active malaria infection at Screening. Testing to be conducted at local laboratories in malariaendemic regions at the discretion of the Investigator.

11. Participant has any confirmed clinically significant drug allergy and/or known hypersensitivity to monoclonal antibody therapeutics or formulation components of the study drug or a related drug.

12. Participant has been in another investigational trial within 30 days or 5 half-lives of the investigational agent (whichever is greater) prior to the Screening Visit.

13. Participant has had a major surgery within 8 weeks prior to the Screening Visit.

14. Participant is pregnant, breastfeeding, or planning to become pregnant during the 48-week treatment period.

15. Participant, parent, or legal guardian are unlikely to comply with the study procedures.

16. Participant has other medical, psychological, or addictive condition that, in the opinion of

the Investigator, would confound or interfere with evaluation of safety, efficacy, and/or PK of the investigational drug; prevent compliance with the study protocol; preclude informed consent; or render the participant, parent, or caretaker unable/unlikely to comply with the study procedures.

Type of study

Interventional

Type of intervention	Type of intervention: Specify t	уре
Pharmaceutical	N/A	
Trial scope	Trial scope: Specify scope	
Safety	N/A	
Study design: Allocation	Study design: Masking	
Randomized controlled trial	Blinded (masking used)	
Study design: Control	Study phase	
Placebo	3	
Study design: Purpose	Study design: Specify purpose	
Treatment	N/A	
Study design: Assignment	Study design: Specify assignr	nent
Parallel	N/A	
IMP has market authorization	IMP has market authorization:	Specify
No		
Name of IMP	Year of authorization	Month of authorization
Inclacumab		
Type of IMP		
Immunological		

Pharmaceutical class



Inclacumab is a recombinant human monoclonal antibody (huMAb) of the immunoglobulin (Ig)G4 subclass directed against human P-selectin, which is being developed by Global Blood Therapeutics,

MINISTRY OF PUBLIC HEALTH

REPUBLIC OF LEBANON Lebanon Clinical Trials Registry

Inc. (GBT), for the treatment of sickle cell disease (SCD). Inclacumab binds to P-selectin, which is a cell adhesion molecule produced by endothelial cells and platelets. Upon activation of these cells (e.g., by thrombin, cytokines, complement components, hypoxia, and heme), P-selectin is translocated to the cell surface where it binds to its primary ligand P-selectin glycoprotein ligand-1 (PSGL-1) and mediates leukocytes recruitment by platelets or endothelial cells. The same mechanism is also responsible for abnormal rolling and adhesion of sickle red blood cells (RBC) to the endothelium, initiating acute vascular occlusion and chronically impairing microvascular blood flow in patients with SCD. Inclacumab binding of P-selectin and prevention of P-selectin binding to its ligands is the putative mechanism by which inclacumab prevents the binding of sickle RBCs or leukocytes to endothelium.

Therapeutic indication

sickle cell disease (SCD)

Therapeutic benefit

Inclacumab is a recombinant huMAb of the IgG4 subclass directed against human P-selectin. The molecule is composed of two

heterodimers, each composed of a heavy and a light polypeptide chain. The four polypeptide chains are linked together by disulfide bonds.

To avoid antibody-dependent cell-mediated cytotoxicity and to improve structural stability, two single point mutations (L235E, S228P) were introduced into the Fc part of the molecule. The inclacumab drug substance is manufactured by fermentation cell culture using Chinese

hamster ovary (CHO) cells followed by purification. The drug substance, drug product, and placebo are manufactured in accordance with Good Manufacturing Practices (GMP).

Results from the SUSTAIN trial in patients with SCD showed that treatment with crizanlizumab, a humanized antibody to P-selectin, resulted in a significantly lower rate of sickle cell-related pain crises (i.e., VOC) than placebo. These data validated P-selectin as a therapeutic target for SCD disease. Inclacumab is currently not approved by any health authority for the treatment of patients with any disease. Inclacumab is being developed to reduce the risk of vaso-occlusive crises in patients with SCD.

Study model Study model: Explain model N/A N/A Study model: Specify model N/A **Time perspective** Time perspective: Explain time perspective N/A N/A Time perspective: Specify perspective N/A Target follow-up duration Target follow-up duration: Unit Number of groups/cohorts **Biospecimen retention Biospecimen description** Samples with DNA** optional genomic samples will be retained beyond study completion



4

REPUBLIC OF LEBANON Lebanon Clinical Trials Registry

Target sample size 240	Actual enrollment target size
Date of first enrollment: Type Anticipated	Date of first enrollment: Date 30/06/2021
Date of study closure: Type Anticipated	Date of study closure: Date 31/10/2023
Recruitment status Pending	Recruitment status: Specify
Date of completion 31/01/2024	
IPD sharing statement plan Yes	IPD sharing statement description Patient's full identity will not be on any of the study documents or sample collected and kept by the sponsor for their studies. Only the partial date of birth will be only collected. Only a unique participant number for the study will link the data or samples to the patients.
Additional data URL	
Admin comments	
Trial status	

. .

Арр	rov	/ea	

Secondary Identifying Numbers		
Full name of issuing authority	Secondary identifying number	
N.A	N.A	

Sources of Monetary or Material Support

Name

Global Blood Therapeutics, Inc.

Secondary Sponsors	
Name	
N.A	





Contact for Public/Scientific Queries						
Contact type	Contact full name	Address	Country	Telephone	Email	Affiliation
Public	Aziz Zoghbi	MCT-CRO, Berytech Technology and Health, 5th Floor Damascus Road, Beirut, Lebanon	Lebanon	009617100 8269	aziz.zoghbi@mct -cro.com	Director of Country Oversight and Manageme nt MENA, Gulf and Africa
Scientific	Miguel Abboud	American University of Beirut Medical Center, Cairo Street, Hamra, Beirut, Lebanon	Lebanon	009613534 213	ma56@aub.edu.l b	PI
Scientific	Adlette Inati	Nini Hospital, el Maarad Street, Triploli, Lebanon	Lebanon	009613228 033	adlette.inati@lau. edu.lb	PI
Scientific	Carolyn Hoppe	181 Oyster Point Blvd. South San Francisco, CA 94080, USA	United States of America	+1 510.289.90 97	choppe@gbt.co m	Medical Monitor

Centers/Hospitals Involved in the Study					
Center/Hospital name	Name of principles investigator	Principles investigator speciality	Ethical approval		
Nini Hospital	Adlette Inati	Professor of Pediatric Hematology and Oncology	Approved		
American University of Beirut Medical Center	Miguel Abboud	Professor of Pediatric Hematology and Oncology	Pending		

Ethics Review				
Ethics approval obtained	Approval date	Contact name	Contact email	Contact phone
Nini Hospital	20/04/2021	Nabil Kabbara	Nabil.kabbara@hopitalnini.com	+961 (0) 6 431 400 ext 1062

Countries of Recruitment	
Name	
Lebanon	

Health Conditions or Problems Studied			
Condition	Code	Keyword	
Sickle Cell Disease	Sickle-cell disorders (D57)	SCD, vaso-occlusive crisis, blood disorders, hemoglobin, red blood cells, sickle-like shape, mutation in hemoglobin gene, sickle-cell trait, sickle-cell crisis, Sickle Cell Disease SCD, Hydroxyurea/ Hydroxycarbamide Therapy, SCA	



Interventions			
Intervention	Description	Keyword	
Inclacumab	Eligible participants will be randomized with a 1:1 ratio into one of two treatment arms as follows: Inclacumab 30 mg/kg administered IV every 12 weeks (Day 1, Week 12, Week 24, and Week 36); or Placebo administered IV every 12 weeks (Day 1, Week 12, Week 24, and Week 36)	Treatment	

Primary Outcomes

Name	Time Points	Measure
Rate of VOCs.	the rate of VOCs during the 48-week treatment period will be compared between the inclacumab and placebo arms with the use of negative binomial regression model.	The incidence of VOC events will be collected every 4 weeks. Each month on non-treatment visit days, participants will be contacted by phone to determine if a VOC event or a pain crisis leading to contact with a healthcare provider without a visit to a medical facility has occurred, to collect AEs, and to record changes to concomitant medications. A VOC is defined as an acute episode of pain that: 1) Has no medically determined cause other than a vaso-occlusive event, and 2) Results in a visit to a medical facility (hospitalization, emergency department, urgent care center, outpatient clinic, or infusion center), or results in a remote contact with a healthcare provider; and 3) Requires parenteral narcotic agents, parenteral nonsteroidal anti-inflammatory drugs (NSAIDs), or an increase in treatment with oral narcotics.
Change in Laboratory assessments.	at each visit	Complete blood count, chemistry and coagulation

Key Secondary Outcomes			
Name	Time Points	Measure	
Time to first VOC during the 48-week treatment period.	48-week treatment period	The secondary efficacy endpoints are based on collection of the incidence and timing of VOCs (per the definition in the primary endpoint) after randomization, QOL assessments and on hospital durations.	
Time to second VOC during the 48-week treatment period.	48-week treatment period	The secondary efficacy endpoints are based on collection of the incidence and timing of VOCs (per the definition in the primary endpoint) after randomization, QOL assessments and on hospital durations.	
Proportion of participants with no VOCs during the 48-week treatment period.	48-week treatment period	The secondary efficacy endpoints are based on collection of the incidence and timing of VOCs (per the definition in the primary endpoint) after randomization, QOL assessments and on hospital durations.	
Rate of VOCs that required admission to a healthcare facility and treatment with parenteral pain medication during the 48- week treatment.	48-week treatment period	The secondary efficacy endpoints are based on collection of the incidence and timing of VOCs (per the definition in the primary endpoint) after randomization, QOL assessments and on hospital durations.	
Number of days of inpatient hospitalization for a VOC during the 48-week treatment period.	48-week treatment period	The secondary efficacy endpoints are based on collection of the incidence and timing of VOCs (per the definition in the primary endpoint) after randomization, QOL assessments and on hospital durations.	
Incidence of treatment-emergent adverse events.	At each visit	Treatment-emergent adverse events (TEAEs).	



Lebanon Clinical Trials Registry

Trial Results	
Summary results	
Study results globally	
Date of posting of results summaries	Date of first journal publication of results
Results URL link	
Baseline characteristics	
Participant flow	
Adverse events	
Outcome measures	
URL to protocol files	